

Supplementary table. Development times, in days, of some Coleoptera species found during the experiment.

Species	T°C	RH	E	L	P	E→A	
<i>Dermestes undulatus</i>	22°	50-60%	5-6	46-60	11	62-76	(Veer et al. 1996)
<i>Dermestes frischii</i>			2-8	22-50	5-12	29-65	(Laudani 1961)
	30°	90-45%	2-4	20-42	6	28-51	(Amos 1968)
	25°	90-45%	3-5	40-86	13	57-105	(Amos 1968)
<i>Creophilus maxillosus</i>			3	16-20 ^a	9-11	30-33	(Kramer 1955)
<i>Aleochara bilineata</i>	22°		5	12-14	8-12	25-31	(Whistlecraft et al.1985)
<i>Aleochara tristis</i>			5-6	6-14	21 ^b	40 ^b	(Drea 1966)
<i>Carcinops pumilio</i>	25,5°	70-80%	5-8	12-19	15-20	34-46	(Morgan et al. 1983)
<i>Necrobia rufipes</i>	30°	80%	4	27-37	9-10	40-51	(Bhuiyan and Saifullah 1997)

T: rearing temperature; RH: relative humidity; E: egg; A: adult; L: larvae; P: pupa; ^a mean time; ^b 1st stage, 2 days; 2nd stage, 2-3 days; 3rd stage, 12-15 days; Gerald (1996) quantifies the length of the 3rd stage at 28°C in 10-14 days; Voris (1939) suggests an incubation of 3 days while the pupa lasts 13 days.

References

- Amos TG 1968. Some laboratory observations on the rates of development, mortality and oviposition of *Dermestes frischii* (Kug.) (Col, Dermestidae). Journal of Stored Products Research 4: 103–117
- Bhuiyan AI, Saifullah AS 1997. Biological note on *Necrobia rufipes* (deg.) (Col.: Cleridae). Bangladesh Journal of Zoology 25: 121–124
- Drea JJ 1966. Studies of *Aleochara tristis* (Coleoptera: Staphylinidae), a natural enemy of the face fly. Journal of Economic Entomology 59: 1368–1373
- Gerald GL 1996. Rearing Techniques for *Creophilus maxillosus*, a Predator of Fly Larvae in Cattle Feedlots. Journal of Economic Entomology 89: 848–851
- Laudani H 1961. Biology and habits of dermestids. Pest Control 29: 58–61
- Morgan P.B, Patterson RS, Weidhaas DE 1983.-A life-history study of *Carcinops pumilio* Erichson (Coleoptera: Histeridae). Journal of the Georgia Entomological Society 18: 353–359
- Veer V, Negi BK, Rao KM, 1996. Dermestid beetles and some other insect pests associated with stored silkworm cocoons in India, including a world list of dermestid species found attacking this commodity. Journal of Stored Products Research 32: 69–89
- Voris R 1939. The Immature Stages of the Genera *Ontholestes*, *Creophilus* and *Staphylinus*. Annals of the Entomological Society of America 32: 289–300
- Whistlecraft JW, Harris CR, Tolman JH, Tomlin AD 1985. Mass-rearing technique for *Aleochara bilineata* (Coleoptera: Staphylinidae). Journal of Economic Entomology 78: 995–997